

Spherical means with centers on a hyperplane in even dimensions

E K Narayanan
Department of Mathematics
Indian Institute of Science
Bangalore, India - 560012

Email: naru@math.iisc.ernet.in

Rakesh
Department of Mathematical Sciences
University of Delaware
Newark, DE 19716, USA

Email: rakesh@math.udel.edu

January 27, 2010

Key words. spherical mean values, wave equation

AMS subject classifications. 35L05, 35L15, 35R30, 44A05, 44A12, 92C55

Abstract

Given a real valued function on $\mathbb{R}^n$ we study the problem of recovering the function from its spherical means over spheres centered on a hyperplane. An old paper of Bukhgeim and Kardakov derived an inversion formula for the odd n case with great simplicity and economy. We apply their method to derive an inversion formula for the even n case. A feature of our inversion formula, for the even n case, is that it does not require the Fourier transform of the mean values or the use of the Hilbert transform, unlike the previously known inversion formulas for the even n case. Along the way, we extend the isometry identity of Bukhgeim and Kardakov for odd n , for solutions of the wave equation, to the even n case.

1 Introduction

Below $n > 1$ will be a positive integer, a point $x \in \mathbb{R}^n$ will occasionally be written as $x = (x', x_n)$ with $x' \in \mathbb{R}^{n-1}$, $x_n \in \mathbb{R}$, and $|S^{n-1}| = \frac{2\pi^{n/2}}{\Gamma(n/2)}$ will be the surface area of the unit sphere in $\mathbb{R}^n$. We take $\hat{f}(\xi) = \int_{\mathbb{R}^n} e^{-ix \cdot \xi} f(x) dx$ as the definition of the Fourier transform of f . Define $\tilde{C}_e^\infty(\mathbb{R}^n)$ to be

the set of smooth functions $f(x)$ on $\mathbb{R}^n$ which vanish¹ to infinite order on $x_n = 0$ and are even in x_n . Let $\tilde{C}_{0e}^\infty(\mathbb{R}^n)$ denote the subspace of compactly supported functions in $\tilde{C}_e^\infty(\mathbb{R}^n)$. Finally, define $L_{e,w}^2(\mathbb{R}^n)$ to be the Hilbert space of measurable functions $f(x)$ on $\mathbb{R}^n$ which are even in x_n and for which $\int_{\mathbb{R}^n} \frac{|f(x)|^2}{|x_n|} dx$ is finite - we use the associated inner product.

For any continuous function $f(x)$ on $\mathbb{R}^n$ define the spherical mean value operator

$$(Mf)(x, t) = \frac{1}{|S^{n-1}|} \int_{|\theta|=1} f(x + t\theta) d\theta, \quad (x, t) \in \mathbb{R}^n \times \mathbb{R}; \quad (1.1)$$

$(Mf)(x, t)$ is the spherical average of f on the sphere centered at x , radius t . The goal is the recovery of f , given $(Mf)(x, t)$ for all x on the hyperplane $x_n = 0$ and for all $t \in \mathbb{R}$. Clearly on $x_n = 0$, $(Mf)(x, t)$ does not change if $f(x)$ is replaced by $f(x', -x_n)$, its reflection across $x_n = 0$. Hence one can recover at most the even, in x_n , part of f . If $f \in \tilde{C}_{0e}^\infty(\mathbb{R}^n)$ then $(Mf)(x, t)$ is an even function of x_n and t and $(Mf)(x', 0, t)$ vanishes to infinite order on $t = 0$. So we define the map $\mathcal{M} : \tilde{C}_{0e}^\infty(\mathbb{R}^n) \rightarrow \tilde{C}_e^\infty(\mathbb{R}^n)$ by

$$(\mathcal{M}f)(x', t) := (Mf)(x', 0, t), \quad (x', t) \in \mathbb{R}^n.$$

The goal of this article is to construct an inverse of $\mathcal{M}$ and characterize the range of $\mathcal{M}$. We will do this indirectly.

There is a well known relationship between spherical mean values and solutions of the wave equation. For any $f \in \tilde{C}_{0e}^\infty(\mathbb{R}^n)$, let $u(x, t)$ be the solution of the initial value problem

$$u_{tt} - \Delta u = 0, \quad \text{on } \mathbb{R}^n \times \mathbb{R}, \quad (1.2)$$

$$u(x, 0) = f(x), \quad u_t(x, 0) = 0, \quad x \in \mathbb{R}^n. \quad (1.3)$$

Note that $u(x, t)$ is even in t and also in x_n because $f(x)$ is even in x_n . From page 682 of [CH62] there is a formula² for $u(x, t)$ in terms of Mf and in terms of f . For any smooth even function $h(t)$ on $\mathbb{R}$ define the differential operator

$$(Dh)(t) := \frac{1}{2t} \frac{\partial h}{\partial t}, \quad t \in \mathbb{R}.$$

Note that $Dh(t)$ makes sense for all t because h is even and $(Dh)(t)$ is an even function of t . Further, D may be considered as differentiation with respect to t^2 because it may be verified that $D(h(t^2)) = h'(t^2)$. We have the following relations for all $(x, t) \in \mathbb{R}^{n+1}$, $t \neq 0$: for odd n

$$u(x, t) = \frac{\sqrt{\pi}}{\Gamma(n/2)} t D^{(n-1)/2} (t^{n-2} (Mf)(x, t)) \quad (1.4)$$

$$= \pi^{(1-n)/2} t D^{(n-1)/2} \int_{\mathbb{R}^n} \delta(t^2 - |x - y|^2) f(y) dy, \quad (1.5)$$

¹ that is $(\partial^\alpha f)(x) = 0$ on $x_n = 0$ for all multi-indices α

²there seems to be a misprint in the formula in [CH62] for the even n case

and for even n

$$u(x, t) = \frac{2}{\Gamma(n/2)} t D^{n/2} \int_0^t \frac{r^{n-1}}{\sqrt{t^2 - r^2}} (Mf)(x, r) dr \quad (1.6)$$

$$= \pi^{-n/2} t D^{n/2} \int_{\mathbb{R}^n} \frac{H(t^2 - |x - y|^2)}{\sqrt{t^2 - |x - y|^2}} f(y) dy \quad (1.7)$$

where $H(s)$ is the Heaviside function. Using (1.2) and that f vanishes to infinite order on $x_n = 0$, it may be verified that $u(x', 0, t)$ vanishes to infinite order on $t = 0$. Define the map $\mathcal{U} : \tilde{C}_{0e}^\infty(\mathbb{R}^n) \rightarrow \tilde{C}_e^\infty(\mathbb{R}^n)$ by

$$(\mathcal{U}f)(x', t) := u(x', 0, t).$$

We construct the inverse of $\mathcal{U}$ and characterize the range of $\mathcal{U}$ which will give us the inverse of $\mathcal{M}$ and implicitly characterize the range of $\mathcal{M}$.

Define the tempered distribution $K(t) := t^{-1/2} H(t)$, $t \in \mathbb{R}$ and let $K^{(m)}(t)$ denote the m -th derivative of $K(t)$.

Theorem 1. *Suppose $n > 1$ is even.*

(a) (Isometry) *For any $f \in \tilde{C}_{0e}^\infty(\mathbb{R}^n)$ we have the isometry*

$$\int_{\mathbb{R}^n} \frac{|f(x)|^2}{|x_n|} dx = \int_{\mathbb{R}^n} \frac{|(\mathcal{U}f)(x', t)|^2}{|t|} dx' dt, \quad (1.8)$$

so $\mathcal{U}$ extends to an isometry on $L_{e,w}^2(\mathbb{R}^n)$.

(b) (Inversion Formula) *For any $f \in \tilde{C}_{0e}^\infty(\mathbb{R}^n)$, $x \in \mathbb{R}^n$, $x_n > 0$ we have*

$$f(x) = \frac{(-1)^{\frac{n}{2}} 2x_n}{\pi^{n/2}} \int_{\mathbb{R}^{n-1}} \int_0^\infty \frac{H(t^2 - x_n^2 - |y'|^2)}{\sqrt{t^2 - x_n^2 - |y'|^2}} t \left(\frac{1}{2t} \frac{\partial}{\partial t} \right)^{n/2} \frac{(\mathcal{U}f)(x' + y', t)}{t} dt dy'. \quad (1.9)$$

(c) (Range Characterization) *The range of $\mathcal{U}$ is a closed subset of $L_{e,w}^2(\mathbb{R}^n)$. Further, if $F \in L_{e,w}^2(\mathbb{R}^n)$ is in the range of the restricted $\mathcal{U}$ (domain is $\tilde{C}_{0e}^\infty(\mathbb{R}^n)$) then*

(i) $F(x', t) \in \tilde{C}_e^\infty(\mathbb{R}^n)$ and $H(\tau) F(x', \sqrt{\tau})/\sqrt{\tau} \in H^s(\mathbb{R}^n)$ for all real s ,

(ii) $|\partial_{x'}^\alpha \partial_t^j F(x', t)| \leq C_{j,\alpha} (1 + |t|)^{-(n-1)/2}$ for all $j \geq 0$ and all multi-indices α .

Finally, if F satisfies (i), (ii) then F is in the range of $\mathcal{U}$ iff $\forall x \in \mathbb{R}^n$, $x_n \neq 0$, we have

$$\int_{\mathbb{R}^{n-1}} \int_0^\infty \frac{H(t^2 + x_n^2 - |y'|^2)}{\sqrt{t^2 + x_n^2 - |y'|^2}} t \left(\frac{1}{2t} \frac{\partial}{\partial t} \right)^{n/2} \frac{F(x' + y', t)}{t} dt dy' = 0. \quad (1.10)$$

In the statement of Theorem 1 we were careful to write only absolutely convergent integrals which forced us into a long-winded statement of (c). A briefer version of (b) and (c) would be that for all $f \in L_{e,w}^2(\mathbb{R}^n)$

$$f(x) = \frac{x_n}{\pi^{n/2}} \int_{\mathbb{R}^n} K^{(n/2)}(t^2 - x_n^2 - |y'|^2) (\mathcal{U}f)(x' + y', t) dy' dt, \quad \forall x \in \mathbb{R}^n, x_n > 0, \quad (1.11)$$

and an $F \in L^2_{e,w}(\mathbb{R}^n)$ is in the range of $\mathcal{U}$ iff

$$\int_{\mathbb{R}^n} K^{(n/2)}(t^2 + x_n^2 - |y'|^2) F(x' + y', t) dy' dt = 0, \quad \forall x \in \mathbb{R}^n, x_n \neq 0. \quad (1.12)$$

However, now (1.11) and (1.12) have to be interpreted in a sense buried in the proof of Theorem 1. The result for odd n given in [BK78] has a form similar to the formal expressions above. They can be rewritten in terms of absolutely convergent integrals similar to the expressions in Theorem 1.

From the hypothesis of Theorem 1 it would seem that the inversion formula is not applicable to an $f \in C_0^\infty(\mathbb{R}^n)$ which is even in x_n but which does not have the right decay near $x_n = 0$. However, for such f , if $u(x', 0, t)$ is known then using $u_{x_n}(x', 0, t) = 0$ (because u is even in x_n) and (1.2) one may determine $f(x', 0) = u(x', 0, 0)$ and all the x_n derivatives of $f(x) = u(x, 0)$ on $x_n = 0$. Knowing these derivatives of $f(x)$ on $x_n = 0$ we can create a known function $g(x) \in C_0^\infty(\mathbb{R}^n)$ so that $\tilde{f} = f - g$ vanishes to a chosen finite order (or infinite order) on $x_n = 0$. Since g is known we can solve (1.2), (1.3) with g replacing f and hence from $u(x', 0, t)$ we can obtain the trace on $x_n = 0$ of the solution of (1.2), (1.3) with f replaced by $\tilde{f}$. Now we can apply the inversion formula to the $\tilde{f}$ data and recover $\tilde{f}$ and hence f .

The early work on the inversion of $\mathcal{U}$ (and hence of $\mathcal{M}$) is described in [Jo55]. For odd n , a nice inversion formula for $\mathcal{U}$ (and hence $\mathcal{M}$) was given by Bukhgeim and Kardakov [BK78]. Later, inversion formulas for $\mathcal{M}$ for odd and even n were derived in [No80], [Fa85], [An88], [NRT95], [Kl03], and [Be09]; [SQ05] has a numerical inversion scheme. The problem under consideration has applications to inverse problems in elasticity and other areas - see [BK78] and [Fa85]. The article [FPR04] studies the similar problem of recovering a compactly supported function from its spherical averages over spheres centered on the boundary of a region containing the support of the function.

Beltukov's formula in [Be09], for the odd n case, is essentially the inversion formula in [BK78]. Beltukov rediscovered some of the ideas in [BK78] (he seems not to have been aware of [BK78]) but not all the ideas in [BK78]. Applying the ideas in [BK78] to the even n case, we give a concise derivation of an inversion formula for the even n case which does not require the Fourier transform of the data or the use of the Hilbert transform. In [BK78] an isometry identity was established for solutions of the wave equation for the odd n case; we extend that identity to the even n case.

The proof of Theorem 1 uses the tempered distribution $N(x) := \pi^{-n/2} K^{(n/2)}(x_n - |x'|^2)$ and its Fourier transform.

Proposition 2. *For even $n > 1$ we have $\widehat{N}(\xi) = e^{i|\xi'|^2/(4\xi_n)}$ for all $\xi \in \mathbb{R}^n$.*

To prove (c) in Theorem 1 and to be sure that the integral in (b) is absolutely convergent we need some decay estimates for solutions of the wave equation - see Theorem 1.1 in [So08].

Proposition 3. Suppose $n > 1$ is a positive integer, $f, g \in C_0^\infty(\mathbb{R}^n)$ and $v(x, t)$ the solution of

$$v_{tt} - \Delta v = 0, \quad \text{on } \mathbb{R}^n \times \mathbb{R}, \quad (1.13)$$

$$v(x, 0) = f(x), \quad v_t(x, 0) = g(x), \quad x \in \mathbb{R}^n \quad (1.14)$$

then

$$\begin{aligned} |v(x, t)| &\lesssim (1 + |t|)^{-(n-1)/2}, & n \text{ odd} \\ |v(x, t)| &\lesssim (1 + |t|)^{-(n-1)/2} (1 + ||t| - |x||)^{-(n-1)/2}, & n \text{ even} \end{aligned} \quad (1.15)$$

with the constant determined by f, g, n .

For n odd, $v(x, t)$ is supported in $|t - |x|| \leq R$ if f, g are supported in $|x| \leq R$. This is not true for n even, instead we have an extra decay contribution over the region $|t - |x|| \geq R$. We have included the statement and the proof for the odd n case because of its possible application to the inversion formula for odd n in [BK78].

The rest of the article is as follows: Section 2 contains the proof of Theorem 1, Section 3 contains the proof of Proposition 2, and Section 4 contains the proof of Proposition 3. We have included the proof of Proposition 3 for completeness though it is stated as a theorem in [So08] whose proof was left as a good exercise. The proof of Theorem 1 would be quite short (see [BK78] for the odd n case) if we accept formal arguments and expressions; our proof is somewhat long because of the need to show the absolute convergence of the integrals in the expressions.

This work was done while the first named author was supported in part by a grant from the UGC via DSA-SAP and the second named author was on sabbatical at the Department of Mathematics of the Indian Institute of Science in Bangalore, India. The second named author thanks the department for its warm hospitality and generous financial support and the University of Delaware for granting the sabbatical.

2 Proof of Theorem 1

For $t > 0$ and $x' \in \mathbb{R}^{n-1}$ from (1.7) we have (integrals below to be interpreted as distributional action)

$$\begin{aligned} \frac{(\mathcal{U}f)(x', 0, t)}{t} &= \frac{u(x', 0, t)}{t} = \pi^{-n/2} \int_{\mathbb{R}^n} K^{(n/2)}(t^2 - |x' - y'|^2 - y_n^2) f(y) dy \\ &= 2\pi^{-n/2} \int_{\mathbb{R}^{n-1}} \int_0^\infty K^{(n/2)}(t^2 - |x' - y'|^2 - y_n^2) f(y) dy_n dy' \\ &= \pi^{-n/2} \int_{\mathbb{R}^{n-1}} \int_0^\infty K^{(n/2)}(t^2 - |x' - q'|^2 - q_n) \frac{f(q', \sqrt{q_n})}{\sqrt{q_n}} dq_n dq'. \end{aligned}$$

Hence taking $x' = p'$, $t = \sqrt{p_n}$ for any $p' \in \mathbb{R}^{n-1}$, $p_n > 0$, we have

$$\frac{(\mathcal{U}f)(p', \sqrt{p_n})}{\sqrt{p_n}} = \pi^{-n/2} \int_{\mathbb{R}^{n-1}} \int_0^\infty K^{(n/2)}(p_n - |p' - q'|^2 - q_n) \frac{f(q', \sqrt{q_n})}{\sqrt{q_n}} dq_n dq'. \quad (2.1)$$

Let $L_+^2(\mathbb{R}^n)$ denote the subspace of functions $g(x)$ in $L^2(\mathbb{R}^n)$ which are zero for $x_n \leq 0$. Define the specialized zero conversion operator $\mathcal{Z} : L_{e,w}^2(\mathbb{R}^n) \rightarrow L_+^2(\mathbb{R}^n)$ with

$$(\mathcal{Z}f)(p) = \begin{cases} \frac{f(p', \sqrt{p_n})}{\sqrt{p_n}} & p_n > 0, \\ 0 & p_n \leq 0. \end{cases}$$

Then $\mathcal{Z}$ is an isomorphism from $L_{e,w}^2(\mathbb{R}^n)$ to $L_+^2(\mathbb{R}^n)$ and its inverse is given by $(\mathcal{Z}^{-1}g)(x', x_n) = x_n^2 g(x', x_n^2)$. Further $\mathcal{Z}$ maps $\tilde{C}_{0e}^\infty(\mathbb{R}^n)$ to the subspace of functions in $C_0^\infty(\mathbb{R}^n)$ which are zero on $x_n \leq 0$.

For any $f \in \tilde{C}_{0e}^\infty(\mathbb{R}^n)$ let $g = \mathcal{Z}f$; then from (2.1) we have

$$(\mathcal{Z}\mathcal{U}f)(p) = \pi^{-n/2} \int_{\mathbb{R}^n} K^{(n/2)}(p_n - |p' - q'|^2 - q_n) g(q) dq = (N * g)(p), \quad p \in \mathbb{R}^n; \quad (2.2)$$

note that this is true even if $p_n \leq 0$. This suggests we define a map $\mathcal{N} : C_0^\infty(\mathbb{R}^n) \rightarrow C^\infty(\mathbb{R}^n)$ with

$$(\mathcal{N}g)(p) = (N * g)(p), \quad p \in \mathbb{R}^n;$$

then $\mathcal{Z}\mathcal{U}f = \mathcal{N}\mathcal{Z}f$. Using Proposition 2 and the definition of $\mathcal{N}$ we have

$$\widehat{\mathcal{N}g}(\xi) = \widehat{N}(\xi) \widehat{g}(\xi) = e^{i|\xi'|^2/(4\xi_n)} \widehat{g}(\xi). \quad (2.3)$$

So $|\widehat{(\mathcal{N}g)}(\xi)| = |\widehat{g}(\xi)|$ for all $\xi \in \mathbb{R}^n$ and $\mathcal{N}$ extends to an isometric isomorphism from $L^2(\mathbb{R}^n)$ to $L^2(\mathbb{R}^n)$. In particular, for all $f \in \tilde{C}_{0e}^\infty(\mathbb{R}^n)$ we have $\|\mathcal{Z}f\|_{L^2(\mathbb{R}^n)} = \|\mathcal{Z}\mathcal{U}f\|_{L^2(\mathbb{R}^n)}$ and hence

$$\int_{\mathbb{R}^{n-1}} \int_0^\infty \frac{|f(p', \sqrt{p_n})|^2}{p_n} dp_n dp' = \int_{\mathbb{R}^{n-1}} \int_0^\infty \frac{|(\mathcal{U}f)(q', \sqrt{q_n})|^2}{q_n} dq_n dq'.$$

Introducing the change of variables $p' = x'$, $p_n = x_n^2$, and $y' = q'$, $t = \sqrt{q_n}$ on the regions $p_n > 0$ and $q_n > 0$ we obtain (1.8) which proves (a) of Theorem 1.

The equation (2.3) allows us to express any $g \in L^2(\mathbb{R}^n)$ in terms of $\mathcal{N}g$ giving us

$$\widehat{g}(\xi) = e^{-i|\xi'|^2/(4\xi_n)} \widehat{\mathcal{N}g}(\xi) = \widehat{N}(\xi', -\xi_n) \widehat{\mathcal{N}g}(\xi). \quad (2.4)$$

Now, for any $g \in C_0^\infty(\mathbb{R}^n)$, $\widehat{g}(\xi)$ is a rapidly decaying function, and $\widehat{N}(\xi', \xi_n)$ is bounded, so $\widehat{\mathcal{N}g}(\xi)$ is a rapidly decaying function and hence $\mathcal{N}g \in H^m(\mathbb{R}^n)$ for all real m . Further $\widehat{N}(\xi', -\xi_n)$ is bounded

so $N(q', -q_n) \in H^{-m}(\mathbb{R}^n)$ for any real $m > n/2$; pick one m . Then for any $p \in \mathbb{R}^n$, using the Fourier inversion formula and (2.4), we have (below T_p is the translation operator)

$$\begin{aligned} g(p) &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \widehat{g}(\xi) e^{ip \cdot \xi} d\xi = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \widehat{N}(\xi', -\xi_n) \widehat{\mathcal{N}g}(\xi) e^{ip \cdot \xi} d\xi \\ &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \overline{\widehat{N}(-\xi', \xi_n)} \widehat{T_p \mathcal{N}g}(\xi) d\xi = \langle N(-q', q_n), (\mathcal{N}g)(p + q) \rangle \\ &= \langle N(p' - q', q_n - p_n), (\mathcal{N}g)(q) \rangle \end{aligned} \quad (2.5)$$

where the $\langle \cdot, \cdot \rangle$ is to be understood as the action of an element in $H^{-m}(\mathbb{R}^n)$ on an element of $H^m(\mathbb{R}^n)$.

For any $f \in \widetilde{C}_{0e}^\infty(\mathbb{R}^n)$, take $g = \mathcal{Z}f$, then $\mathcal{N}g = \mathcal{Z}\mathcal{U}f$; hence for any $p \in \mathbb{R}^n$, from (2.5) we have

$$(\mathcal{Z}f)(p) = \langle N(p' - q', q_n - p_n), (\mathcal{Z}\mathcal{U}f)(q) \rangle. \quad (2.6)$$

Working formally, for the moment, (2.6) implies

$$\begin{aligned} (\mathcal{Z}f)(p) &= \langle K^{(n/2)}(p' - q', q_n - p_n), (\mathcal{Z}\mathcal{U}f)(q) \rangle \\ &= (-1)^{n/2} \langle K(p' - q', q_n - p_n), \left(\frac{\partial}{\partial q_n} \right)^{n/2} (\mathcal{Z}\mathcal{U}f)(q) \rangle \\ &= (-\pi)^{n/2} \int_{\mathbb{R}^n} \frac{H(q_n - p_n - |q' - p'|^2)}{\sqrt{q_n - p_n - |q' - p'|^2}} \left(\frac{\partial}{\partial q_n} \right)^{n/2} (\mathcal{Z}\mathcal{U}f)(q', q_n) dq. \end{aligned}$$

For any $x \in \mathbb{R}^n$ with $x_n > 0$, taking $p' = x'$, $p_n = x_n^2$, and taking $y' = q'$, $t = \sqrt{q_n}$ (note $q_n > 0$ on the support of $\mathcal{Z}\mathcal{U}f$), we have

$$\frac{f(x)}{x_n} = 2(-\pi)^{-n/2} \int_{\mathbb{R}^{n-1}} \int_0^\infty \frac{H(t^2 - x_n^2 - |x' - y'|^2)}{\sqrt{t^2 - x_n^2 - |x' - y'|^2}} t \left(\frac{1}{2t} \frac{\partial}{\partial t} \right)^{n/2} \frac{(\mathcal{U}f)(y', t)}{t} dt dy'$$

thus proving (b) of Theorem 1.

We now address (c) of Theorem 1. If $f \in \widetilde{C}_{0e}^\infty(\mathbb{R}^n)$ then $(\mathcal{U}f)(x', t) \in \widetilde{C}_e^\infty(\mathbb{R}^n)$ and from (2.3) $\mathcal{Z}\mathcal{U}f \in H^s(\mathbb{R}^n)$ for all real s . Further, if u is the solution of (1.2), (1.3) then $\partial_x^\alpha \partial_t^m u$ is a solution of (1.13), (1.14) for some $f, g \in C_0^\infty(\mathbb{R}^n)$ and hence from Proposition 3

$$|\partial_x^\alpha \partial_t^m (\mathcal{U}f)(x', t)| \leq C_{m,\alpha} (1 + |t|)^{-(n-1)/2}$$

for all $m \geq 0$ and all multi-indices α , proving (i), (ii) of (c) in Theorem 1.

The extended $\mathcal{U}$ is an isometry and hence its range is closed. A function $F(x', t) \in L_{e,w}^2(\mathbb{R}^n)$ is in the range of $\mathcal{U}$ iff $F = \mathcal{U}f$ for some $f \in L_{e,w}^2(\mathbb{R}^n)$, that is (using the isomorphism of $\mathcal{Z}$) iff $\mathcal{Z}F = \mathcal{Z}\mathcal{U}f = \mathcal{N}\mathcal{Z}f$, that is (using the isomorphism of $\mathcal{N}$) iff $\mathcal{Z}f = \mathcal{N}^{-1}\mathcal{Z}F$ for some $f \in L_{e,w}^2(\mathbb{R}^n)$. So a necessary condition for F to be in the range of $\mathcal{U}$ is that $(\mathcal{N}^{-1}\mathcal{Z}F)(p) = 0$ for $p_n < 0$. Conversely, if $(\mathcal{N}^{-1}\mathcal{Z}F)(p) = 0$ on the region $p_n < 0$, then $g := \mathcal{N}^{-1}\mathcal{Z}F$ is in $L_+^2(\mathbb{R}^n)$. Let

$f := \mathcal{Z}^{-1}g$; then $f \in L^2_{e,w}(\mathbb{R}^n)$ and $g = \mathcal{Z}f$ and hence $\mathcal{Z}f = \mathcal{N}^{-1}\mathcal{Z}F$ implying $\mathcal{Z}F = \mathcal{N}\mathcal{Z}f = \mathcal{Z}\mathcal{U}f$, so $F = \mathcal{U}f$ proving our claim.

If F satisfies (i), (ii) of (c) in Theorem 1, then $\mathcal{Z}F \in H^s(\mathbb{R}^n)$ for all s . From (2.4), $(\widehat{\mathcal{N}^{-1}\mathcal{Z}F})(\xi) = N(\xi', -\xi_n)\widehat{\mathcal{Z}F}(\xi)$ and since the right side is an integrable function so is the left side, and hence by the Fourier inversion formula, as done for (2.5), we have

$$(\mathcal{N}^{-1}\mathcal{Z}F)(p) = \langle N(p' - q', q_n - p_n), (\mathcal{Z}F)(q) \rangle, \quad p \in \mathbb{R}^n.$$

Hence, working formally for the moment, F is in the range only if for all $p' \in \mathbb{R}^{n-1}$, with $p_n > 0$ we have (as done earlier)

$$\begin{aligned} 0 &= (\mathcal{N}^{-1}\mathcal{Z}F)(p', -p_n) = \langle N(p' - q', q_n + p_n), (\mathcal{Z}F)(q) \rangle \\ &= (-\pi)^{n/2} \int_{\mathbb{R}^n} \frac{H(q_n + p_n - |q' - p'|^2)}{\sqrt{q_n + p_n - |q' - p'|^2}} \left(\frac{\partial}{\partial q_n} \right)^{n/2} (\mathcal{Z}F)(q', q_n) dq \\ &= (-\pi)^{n/2} \int_{\mathbb{R}^{n-1}} \int_0^\infty \frac{H(q_n + p_n - |p' - q'|^2)}{\sqrt{q_n + p_n - |p' - q'|^2}} \left(\frac{\partial}{\partial q_n} \right)^{n/2} \frac{F(q', \sqrt{q_n})}{\sqrt{q_n}} dq_n dq'. \end{aligned}$$

For any $x \in \mathbb{R}^n$ with $x_n \neq 0$, take $p' = x'$, $p_n = x_n^2$. Also, inside the integral use the change of variables $y' = q'$, $t^2 = q_n$. Then F is in the range of $\mathcal{U}$ iff for all $x \in \mathbb{R}^n$ with $x_n \neq 0$ we have

$$0 = \int_{\mathbb{R}^{n-1}} \int_0^\infty \frac{H(t^2 + x_n^2 - |x' - y'|^2)}{\sqrt{t^2 + x_n^2 - |x' - y'|^2}} t \left(\frac{1}{2t} \frac{\partial}{\partial t} \right)^{n/2} \frac{F(y', t)}{t} dt dy'$$

proving (c) of Theorem 1.

So to complete a rigorous proof of the Theorem, for both parts (b) and (c), we have to show that if F satisfies (i), (ii) of part (c) of Theorem 1, then for all $p \in \mathbb{R}^n$, we have

$$\langle N(p' - q', q_n - p_n), (\mathcal{Z}F)(q) \rangle = (-\pi)^{n/2} \int_{\mathbb{R}^n} \frac{H(q_n - p_n - |q'|^2)}{\sqrt{q_n - p_n - |q'|^2}} \left(\frac{\partial}{\partial q_n} \right)^{n/2} (\mathcal{Z}\mathcal{U}f)(q' + p', q_n) dq,$$

where the inner product $\langle \cdot, \cdot \rangle$ is to be interpreted as the action of an element in $H^{-m}(\mathbb{R}^n)$ on an element of $H^m(\mathbb{R}^n)$. We now give a proof of this claim.

For any $\phi \in C_0^\infty(\mathbb{R}^n)$, the value of $\langle N(p' - q', q_n - p_n), \phi(q) \rangle$, as determined by the duality between H^{-m} and H^m , is the same as the value of the action of the tempered distribution $N(p' - q', q_n - p_n)$ on $\phi(q)$. Let $\chi \in C^\infty(\mathbb{R})$ with $\chi(s) = 1$ for $s \leq 0$ and $\chi(s) = 0$ for $s \geq 1$ and $\psi \in C^\infty(\mathbb{R}^{n-1})$ with $\psi(x') = 1$ if $|x'| \leq 1$ and $\psi(x') = 0$ if $|x'| \geq 2$. Then

$$\phi_k(q) := \chi(\log(q_n^2) - k) \psi(q'/k) (\mathcal{Z}F)(q)$$




Figure 1: Graphs of $\phi(s)$ and $\psi(x')$

is in $C_0^\infty(\mathbb{R}^n)$ and $\phi_k(q)$ converges to $(\mathcal{Z}F)(q)$ in $H^m(\mathbb{R}^n)$ for all m . Hence

$$\begin{aligned}
\langle N(p' - q', q_n - p_n), (\mathcal{Z}F)(q) \rangle &= \lim_{k \rightarrow \infty} \langle N(p' - q', q_n - p_n), \phi_k(q) \rangle \\
&= \lim_{k \rightarrow \infty} (-\pi)^{n/2} \langle K(q_n - p_n - |p' - q'|^2), (\partial/\partial q_n)^{n/2} \phi_k(q) \rangle \\
&= \lim_{k \rightarrow \infty} (-\pi)^{n/2} \sum_{j=0}^{n/2} \binom{n/2}{j} \langle K(q_n - p_n - |p' - q'|^2), \partial_n^j (\chi(\log(q_n^2) - k)) \psi(q'/k) \partial_n^{n/2-j} (\mathcal{Z}F)(q) \rangle \\
&= \lim_{k \rightarrow \infty} (-\pi)^{n/2} \langle K(q_n - p_n - |p' - q'|^2), \chi(\log(q_n^2) - k) \psi(q'/k) \partial_n^{n/2} (\mathcal{Z}F)(q) \rangle \tag{2.7}
\end{aligned}$$

$$= (-\pi)^{n/2} \int_{\mathbb{R}^n} \frac{H(q_n - p_n - |q'|^2)}{\sqrt{q_n - p_n - |q'|^2}} \left(\frac{\partial}{\partial q_n} \right)^{n/2} (\mathcal{Z}F)(q' + p', q_n) dq. \tag{2.8}$$

The equations above required the observation that

$$\lim_{k \rightarrow \infty} \langle K(q_n - p_n - |p' - q'|^2), \partial_n^j (\chi(\log(q_n^2) - k)) \psi(q'/k) \partial_n^{n/2-j} (\mathcal{Z}F)(q) \rangle = 0, \quad j > 0 \tag{2.9}$$

and that (2.7) equals (2.8). Both these observations follow from the use of the dominated convergence theorem, that for $j > 0$, $\partial_n^j (\chi(\log(q_n^2) - k))$ is supported in $e^{k/2} \leq |q_n| \leq e^{(k+1)/2}$, and the integrability in q of $K(q_n - p_n - |p' - q'|^2) (1 + |q_n|)^{-j} \partial_n^{n/2-j} (\mathcal{Z}F)(q)$ for $j = 0 \cdots n/2$.

We now prove the integrability assertion in the previous sentence. For any function $h(t)$ and any real number k we have $\frac{1}{2t} \frac{\partial}{\partial t} (t^k h(t)) = c_1 t^{k-2} h(t) + c_2 t^{k-1} \partial_t h(t)$, so for any $m \geq 0$ we have

$$\left(\frac{1}{2t} \frac{\partial}{\partial t} \right)^m (t^k h(t)) = \sum_{l=0}^m c_l t^{k-l-2(m-l)} \partial_t^l h(t) = \sum_{l=0}^m c_l t^{k-2m+l} \partial_t^l h(t). \tag{2.10}$$

Hence using the substitution $q_n = t^2$ we have

$$\left(\frac{\partial}{\partial q_n} \right)^m (\mathcal{Z}F)(q) = \left(\frac{\partial}{\partial q_n} \right)^m \frac{F(q', \sqrt{q_n})}{\sqrt{q_n}} = \left(\frac{1}{2t} \frac{\partial}{\partial t} \right)^m \frac{F(q', t)}{t} = \sum_{l=0}^m t^{-1-2m+l} \partial_t^l F(q', t)$$

and from the decay properties of F we conclude that

$$\left| \left(\frac{\partial}{\partial q_n} \right)^m (\mathcal{Z}F)(q) \right| \lesssim (1 + |t|)^{-1-m} (1 + |t|)^{-(n-1)/2} |_{t=\sqrt{q_n}} \lesssim (1 + |q_n|)^{-(2m+n+1)/4}. \quad (2.11)$$

Hence for any $j \geq 0$,

$$\begin{aligned} & \int_{\mathbb{R}^n} \frac{K(q_n - p_n - |p' - q'|^2)}{(1 + |q_n|)^j} |\partial_n^{(n/2)-j} (\mathcal{Z}F)(q)| dq \\ &= \int_{p_n}^{\infty} \int_{|q'|^2 \leq q_n - p_n} \frac{|\partial_n^{(n/2)-j} (\mathcal{Z}F)(q' + p', q_n)|}{\sqrt{q_n - p_n - |q'|^2}} (1 + |q_n|)^{-j} dq' dq_n \\ &= \int_{p_n}^{\infty} \int_{|z'|^2 \leq 1} \frac{|\left(\partial_n^{(n/2)-j} (\mathcal{Z}F) \right) (\sqrt{q_n - p_n} z' + p', q_n)|}{\sqrt{1 - |z'|^2}} (1 + |q_n|)^{-j} (q_n - p_n)^{(n-2)/2} dz' dq_n \\ &\lesssim \int_{p_n}^{\infty} \int_{|z'|^2 \leq 1} \frac{(1 + |q_n|)^{-5/4}}{\sqrt{1 - |z'|^2}} dz' dq_n \end{aligned}$$

which is finite; we have used (2.11) and noted that for all $j = 0 \cdots n/2$

$$(-j + (n-2)/2) - (n+1 + 2(n/2 - j))/4 = -j/2 - 5/4 \leq -5/4.$$

QED

3 Proof of Proposition 2

Let $\mathcal{S}(\mathbb{R}^n)$ denote the usual space of smooth, rapidly decreasing functions on $\mathbb{R}^n$. Choose $\chi(s) \in C^\infty(\mathbb{R})$ with $\chi(s) = e^{-s}$ for $s \geq 0$ and $\chi(s) = 0$ for $s \leq -1$; then χ_ϵ , defined through

$$\widehat{\chi}_\epsilon(x) := e^{-\epsilon^2 |x'|^2} \chi(\epsilon x_n), \quad x \in \mathbb{R}^n,$$

satisfies $\lim_{\epsilon \rightarrow 0^+} \chi_\epsilon * \phi = \phi$ for all $\phi \in \mathcal{S}(\mathbb{R}^n)$ in the topology of $\mathcal{S}(\mathbb{R}^n)$.

If $P(x) = K(x_n - |x'|^2)$ then $N(x) = \pi^{-n/2} \partial_n^{n/2} P(x)$ and hence $\widehat{N}(\xi) = \pi^{-n/2} (i \xi_n)^{n/2} \widehat{P}(\xi)$. By definition, for all $\phi \in \mathcal{S}(\mathbb{R}^n)$ we have

$$\langle \widehat{P}, \phi \rangle = \lim_{\epsilon \rightarrow 0^+} \langle \widehat{P}, \chi_\epsilon * \phi \rangle = \lim_{\epsilon \rightarrow 0^+} \langle P, \widehat{\chi}_\epsilon \widehat{\phi} \rangle. \quad (3.1)$$

Now (all integrals below are absolutely convergent)

$$\begin{aligned} \langle P, \widehat{\chi}_\epsilon \widehat{\phi} \rangle &= \int_{|x'|^2 \leq x_n} \frac{e^{-\epsilon^2 |x'|^2} \chi(\epsilon x_n)}{\sqrt{x_n - |x'|^2}} \widehat{\phi}(x) dx = \int_{\mathbb{R}^{n-1}} \int_{|x'|^2}^{\infty} \int_{\mathbb{R}^n} \frac{e^{-\epsilon^2 |x'|^2} e^{-\epsilon x_n}}{\sqrt{x_n - |x'|^2}} e^{-ix \cdot \xi} \phi(\xi) d\xi dx_n dx' \\ &= \int_{\mathbb{R}^n} F_\epsilon(\xi) \phi(\xi) d\xi \end{aligned}$$

where

$$\begin{aligned}
F_\epsilon(\xi) &= \int_{\mathbb{R}^{n-1}} \int_{|x'|^2}^\infty \frac{e^{-\epsilon^2|x'|^2} e^{-\epsilon x_n}}{\sqrt{x_n - |x'|^2}} e^{-ix \cdot \xi} dx_n dx' \\
&= \int_{\mathbb{R}^{n-1}} e^{-(\epsilon^2 + \epsilon + i\xi_n)|x'|^2} e^{-ix' \cdot \xi'} dx' \int_0^\infty \frac{e^{-x_n(\epsilon + i\xi_n)}}{\sqrt{x_n}} dx_n \\
&= \frac{\pi^{(n-1)/2}}{(\sqrt{\epsilon^2 + \epsilon + i\xi_n})^{n-1}} \exp\left(\frac{-|\xi'|^2}{4(\epsilon^2 + \epsilon + i\xi_n)}\right) \frac{\sqrt{\pi}}{\sqrt{\epsilon + i\xi_n}}
\end{aligned}$$

and the branch of square root used is the one obtained by cutting out the negative reals and whose argument lies in $(-\pi/2, \pi/2)$. Hence

$$\begin{aligned}
\langle \widehat{N}, \phi \rangle &= \langle \pi^{-n/2} (i\xi_n)^{n/2} \widehat{P}(\xi), \phi(\xi) \rangle = \langle \widehat{P}(\xi), \pi^{-n/2} (i\xi_n)^{n/2} \phi(\xi) \rangle \\
&= \lim_{\epsilon \rightarrow 0^+} \int_{\mathbb{R}^n} \pi^{-n/2} (i\xi_n)^{n/2} F_\epsilon(\xi) \phi(\xi) d\xi.
\end{aligned}$$

Now $\pi^{-n/2} (i\xi_n)^{n/2} F_\epsilon(\xi)$ may be seen to be bounded with a bound independent of ϵ and its pointwise limit is $\exp(i|\xi'|^2/(4\xi_n))$ almost everywhere, hence by the dominated convergence theorem

$$\langle \widehat{N}, \phi \rangle = \int_{\mathbb{R}^n} \exp(i|\xi'|^2/(4\xi_n)) \phi(\xi) d\xi.$$

QED

4 Proof of Proposition 3

Because of linearity it is enough to deal only with the two special cases where either $f = 0$ or $g = 0$. We give the proof only of the case where $g = 0$, in which case v is given by the formulas (1.5), (1.7) for n odd, even respectively. The case where $f = 0$ is dealt with similarly with the formula for the new v being almost the same as (1.5), (1.7) except the t term is dropped, the D power is reduced by one, and the expressions must be multiplied by 2.

4.1 Odd n case

Let $f \in C_0^\infty(\mathbb{R}^n)$ with f supported in the 0 centered ball of radius R . Since $(Mf)(x, t) \neq 0$ iff the sphere $|y - x| = t$ intersects the ball $|y| \leq R$, we see that $(Mf)(x, t)$ is supported in the region $|t - |x|| \leq R$, hence, using (1.4), $u(x, t)$ is supported in the region $|t - |x|| \leq R$. Now suppose

$t > 0$, p a real number and $h \in C_0^\infty(\mathbb{R}^n)$; then

$$\begin{aligned} D(t^p(Mh))(x, t) &= \frac{p}{2} t^{p-2} (Mh)(x, t) + \frac{t^{p-1}}{2|S^{n-1}|} \int_{|\theta|=1} \theta \cdot (\nabla h)(x + t\theta) d\theta \\ &= \frac{p}{2} t^{p-2} (Mh)(x, t) + \frac{t^{p-2}}{2|S^{n-1}|} \int_{|\theta|=1} (x + t\theta - x) \cdot (\nabla h)(x + t\theta) d\theta \\ &= c_1 t^{p-2} (Mh)(x, t) + c_2 t^{p-2} M(y \cdot \nabla h(y))(x, t) + c_3 t^{p-2} x \cdot M(\nabla h)(x, t) \end{aligned} \quad (4.1)$$

$$= t^{p-2} \sum_{|\alpha| \leq 1} x^\alpha M(h_\alpha)(x, t) \quad (4.2)$$

where $h_\alpha(x)$ are smooth functions which are just made of derivatives of h , possibly multiplied with powers of x . Hence, using (4.2) $(n-1)/2$ times in (1.4), we see that

$$u(x, t) = c t D^{(n-1)/2} t^{n-2} (Mf)(x, t) = \sum_{|\alpha| \leq (n-1)/2} x^\alpha (Mh_\alpha)(x, t) \quad (4.3)$$

where h_α are made up of derivatives of f , possibly multiplied with powers of x . Now for any h supported in $|x| \leq R$, the surface area of the intersection of the sphere $|y - x| = t$ with the ball $|y| \leq R$ will be bounded above by a constant independent of x or t . Hence for $t \geq 1$

$$|(Mh)(x, t)| \leq \frac{1}{t^{n-1} |S^{n-1}|} \int_{|y-x|=t} |h(y)| dS_y \lesssim (1+t)^{-(n-1)} \quad (4.4)$$

and noting that u is supported in $|t - |x|| \leq R$, for $t > 1$ we have from (4.3) that

$$|u(x, t)| \lesssim \frac{(1+|x|)^{(n-1)/2}}{(1+t)^{n-1}} \lesssim (1+|t|)^{-(n-1)/2}.$$

4.2 Even n case

First note that $u(x, t)$ is supported in $|x| \leq t + R$. We will show the decay rates, as $t \rightarrow \infty$, for the two cases $|x| - R \leq t \leq |x| + 3R$ and $|x| + 3R \leq t$ separately.

Suppose $|x| + 3R \leq t$; then for any y in the support of f we have $|x - y| \leq |x| + |y| \leq |x| + R \leq t$. Hence the region $|x - y| \leq t$ includes the support of f . So from (1.7) we have

$$\begin{aligned} |u(x, t)| &= \pm c t D^{n/2} \int_{|y| \leq R} \frac{f(y)}{\sqrt{t^2 - |x - y|^2}} dy = \pm c t \int_{|y| \leq R} \frac{f(y)}{(t^2 - |x - y|^2)^{(n+1)/2}} dy \\ &\lesssim \frac{t}{(t^2 - (|x| + R)^2)^{(n+1)/2}} = \frac{t}{(t + |x| + R)^{(n+1)/2}} \frac{1}{(t - |x| - R)^{(n+1)/2}}. \end{aligned}$$

Now for $|x| + 3R \leq t$ we have $R \leq (t - |x|)/3$ and

$$t - |x| - R \geq t - |x| - 2R + R \geq t - |x| - \frac{2}{3}(t - |x|) + R \geq c(|t - |x|| + 1),$$

hence

$$|u(x, t)| \lesssim (1+t)^{-(n-1)/2} (1+|t-|x||)^{-(n+1)/2}.$$

This estimate is better than the estimate in the statement of the proposition - the weaker estimate in the statement of the proposition is tight when $f = 0$ and g is non-zero.

For $|x| - R \leq t \leq |x| + 3R$ we use a technique similar to one used in the odd case. Since we are interested in situation when t is large, we may assume that $|x| > \max(1, R)$. From (1.7) we have

$$u(x, t) = ct D^{n/2} \int_0^t \frac{r^{n-1}}{\sqrt{t^2 - r^2}} (Mf)(x, r) dr = ct D^{n/2} t^{n-1} \int_0^1 \frac{s^{n-1}}{\sqrt{1 - s^2}} (Mf)(x, ts) ds. \quad (4.5)$$

Now, for $t > 0$, for any real number p and any smooth function $h(x)$ we have

$$\begin{aligned} D(t^p (Mh)(x, st)) &= \frac{p}{2} t^{p-2} (Mh)(x, st) + \frac{t^{p-1}}{2|S^{n-1}|} \int_{|\theta|=1} s\theta \cdot (\nabla h)(x + st\theta) d\theta \\ &= \frac{p}{2} t^{p-2} (Mh)(x, st) + \frac{t^{p-2}}{2|S^{n-1}|} \int_{|\theta|=1} (x + st\theta - x) \cdot (\nabla h)(x + st\theta) d\theta \\ &= c_1 t^{p-2} (Mh)(x, st) + c_2 t^{p-2} M(y \cdot \nabla h(y))(x, st) + c_3 t^{p-2} x \cdot M(\nabla h)(x, st) \\ &= t^{p-2} \sum_{|\alpha| \leq 1} x^\alpha M(h_\alpha)(x, st) \end{aligned}$$

where $h_\alpha(x)$ are smooth functions which are just made of derivatives of h , possibly multiplied with powers of x . Hence

$$t D^{n/2} t^{n-1} (Mf)(x, ts) = \sum_{|\alpha| \leq n/2} x^\alpha (Mh_\alpha)(x, st);$$

using this in (4.5), noting that $(Mh_\alpha)(x, st)$ is supported in $|x| - R \leq st \leq |x| + R$, and using (4.4) with h replaced by h_α , we obtain (note $|x| \geq R$)

$$\begin{aligned} |u(x, t)| &\lesssim |x|^{n/2} \int_0^1 \frac{s^{n-1}}{\sqrt{1 - s^2}} |(Mh_\alpha)(x, st)| ds \lesssim \frac{|x|^{n/2}}{t^{n-1}} \int_{(|x|-R)/t}^1 \frac{1}{\sqrt{1 - s^2}} ds \\ &= \frac{|x|^{n/2}}{t^{n-1}} \arccos((|x| - R)/t). \end{aligned}$$

We are considering the region where $t - 3R \leq |x| \leq t + R$, and there is no loss of generality in assuming that $t > \max(1, 4R)$; hence

$$|u(x, t)| \lesssim (1+t)^{-(n-2)/2} \arccos((t - 4R)/t) \lesssim (1+t)^{-(n-1)/2}$$

where the last step follows from $\lim_{t \rightarrow \infty} t^{1/2} \arccos((t - 4R)/t) = c \neq 0$ by L'Hopital's rule. So Proposition 3 follows from the fact that $R \leq |t - |x|| \leq 3R$ in the region of interest. **QED**

References

- [An88] Andersson, Lars-Erik. On the determination of a function from spherical averages. *SIAM J. Math. Anal.* 19 (1988), no. 1, 214–232.
- [Be09] Beltukov, A. Inversion of the spherical mean transform with sources on a hyperplane, *Math Arxiv* arXiv:0910.1380 (2009).
- [BK78] Bukhgeim, A L and Kardakov, V B. Solution of an inverse problem for an elastic wave equation by the method of spherical means. *Siberian Math Journal*, 19 (1978), no. 4, 749–758.
- [CH62] Courant, R and Hilbert, D. *Methods of Mathematical Physics, Volume II*, John Wiley, New York, 1962.
- [Fa85] Fawcett, J A. Inversion of n-dimensional spherical averages, *SIAM J. Appl. Math.*, 45 (1985), 336341.
- [FPR04] Finch, D; Patch, S and Rakesh. Determining a function from its mean values over a family of spheres, *SIAM J. Math. Anal.* 35, No. 5, 1213-1240 (2004).
- [Kl03] Klein, J. Inverting the spherical radon transform for physically meaningful functions, eprint arXiv:math/0307348 (2003).
- [NRT95] Nessibi, M M; Rachdi, L T and Trimeche, K. Ranges and inversion formulas for spherical mean operator and its dual, *J. Math. Anal. Appl.* 196 (1995), no. 3, 861884.
- [No80] Norton, S J. Reconstruction of a reflectivity field from line integrals over circular paths, *J. Acoust. Soc. Amer.* 67 (1980), no. 3, 853863.
- [SQ05] Schuster, T and Quinto, E T. On a regularization scheme for linear operators in distribution spaces with an application to the spherical Radon transform, *SIAM J. Appl. Math.*, 65:1369-1387, (2005).
- [So08] Sogge, C D. *Lectures on non-linear wave equations*, Second Edition, International Press (2008).